

What is Plenvu

Each box contains 3 sachets, making up dose 1 (1 sachet), and dose 2 (sachet A & B)

Why do I need to take it?

Taking Plenvu before your appointment will give you a clean bowel.

What to Expect

After you drink Plenvu you will have lots of watery bowel movements, so stay near a toilet because you may need to use it urgently.



Reporting of Side Effects

United Kingdom

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

Ireland

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly to the HPRa, via www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

A patient support item from Norgine.

Norgine Pharmaceuticals Limited, Moorhall Road, Harefield, Middlesex UB9 6NS.

E-mail: os-medinfo@norgine.com.

Tel: +44 (0) 1895 826 606.

Importance of Proper Cleansing

- A clean bowel will allow for a clear view of its inner lining which is important for both accurate diagnosis and treatment.
- Abnormal changes may be tiny, so even the smallest amount of faeces could potentially disguise something important.
- **If the bowel is not empty and clean enough, the procedure may have to be cancelled and repeated on another occasion at a later date.**

How to Prepare Plenvu

Dose 1



Empty dose 1 sachet in to a 500ml jug



Add water to make up 500ml and stir until fully dissolved



Sip very slowly over **60 minutes**

Dose 2



Empty dose 2 sachets (A & B) in to a 500ml jug



Add water to make up 500ml and stir until fully dissolved



Sip very slowly over **60 minutes**



drink a minimum of **500ml** clear fluids with each dose.

Alternating between this and the Plenvu solution is acceptable

Patient Advice Clear Fluids

You must drink a minimum of 500ml extra clear fluids with each dose. Continue drinking clear fluids to keep yourself hydrated until 2 hours before your procedure and then stop drinking, clear drinks may include:



Water



Clear soups (no bits)



Tea & Coffee (without milk)



Soft drinks (cordial - not blackcurrant)

••••• Do Not Drink Alcohol •••••

Plenvu may be easier to take when:

- Adding cordial to the additional clear fluids (to alter the flavour, but not blackcurrant as it will stain the bowel)

- It has been chilled once it has been made up

Further advice for patients:

- Using wet wipes after each bowel movement
- Use a barrier cream to prevent a 'sore bottom' for example: petroleum jelly

Hints & Tips



Sip the Plenvu solution **very slowly** over **60 minutes** alongside a minimum of 500ml clear fluids (you can alternate between sips of Plenvu and clear fluid if you'd like)



Once dissolved (this may take up to 8 minutes) the **solution can be refrigerated**



Try using a **straw**



Stay hydrated with plenty of clear fluids

When to Take Your Plenvu

Day Before Dosing

Day Before Procedure				Day Of Procedure		
13:00	17:00	21:00				
Begin fasting	Take 1st Dose	Take 2nd Dose	Sleep	Stop Drinking	Your Appointment	
 Low fibre diet, see dietary advice	 Fasting Period: NO FOOD. Do not eat again until after your procedure	 1st Dose Mix with 500 ml of water +  drink a minimum 500ml clear fluids with each dose Sip slowly over 60 minutes Alternating between Plenvu and clear fluid is acceptable	 2nd Dose Mix with 500 ml of water +  drink a minimum 500ml clear fluids with each dose Sip slowly over 60 minutes Alternating between Plenvu and clear fluid is acceptable	 Sleep	 2 hours before appointment or as advised NO DRINK	 Your Appointment

Dietary Advice

Your healthcare professional may request that you adopt a low fibre diet when preparing for your bowel preparation. For specialised diets, please refer to your healthcare professional


••••• Low Fibre Foods Which Are Preferred •••••

 Butter or margarine	 Cheese, eggs or tofu	 Clear soups, tea or coffee	 Ice cream or custard
 Natural plain yoghurt	 White meats, skinless chicken, grilled or poached fish	 Shredless jam or marmalade	 White bread, rice, pasta, boiled or mashed potatoes

••••• High Fibre Foods Which Should Be Avoided •••••

 Red meat, sausages or pies	 Nuts, pulses, seeds inc. baked beans	 Wholemeal pasta or brown rice	 Vegetables, fruit or salad
 Cakes or biscuits	 Potato skins, chips or french fries	 Puddings containing fruit or nuts	 Wholemeal, seeded bread or brown bread

What is a Colonoscopy?



A colonoscopy is a test that uses a narrow, flexible, telescopic camera called a colonoscope to look at the lining of your large bowel. A colonoscopy can be used to detect irritated and swollen tissue, ulcers, or growths such as polyps. It helps to diagnose or exclude health conditions like Crohn's disease, ulcerative colitis, diverticular disease and bowel cancer.

A patient video and more advice on taking Plenvu can be found by visiting:

bowelcleansingmatters.co.uk

This is an external link to a patient website intended for those patients who have been prescribed Plenvu by their healthcare professional.



Morning Appointment

Dosing timeline code: UKE-COR-PR-2300012
 Date of preparation: July 2023
 What is a Colonoscopy Code : UKE-COR-PR-2200017
 Date of preparation: July 2022
 Code: UKE-COR-PR-2300009
 Date of preparation: July 2023

